

Appl. No. 10/643,681
Docket No.: 18528.642/ 254/057CON
Page 2

AMENDMENTS TO THE CLAIMS

Please enter the following amendments without prejudice or disclaimer.

This listing of claims will replace all prior versions, and listings, of claims in the application.

In the claims:

1-23. (Canceled)

24. (Currently amended) A method of reducing or moderating a postprandial rise in plasma glucose in a mammal comprising administering to said mammal an amylin or an amylin agonist analogue in an amount effective to reduce or moderate a postprandial rise in plasma glucose, wherein the amylin agonist analogue is a peptide.

25. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-
G₁Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-J₁-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or Hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Ser, Pro or Thr;

K₁ is Asn, Asp or Gln;

Appl. No. 10/643,681

Docket No.: 18528.642/ 254/057CON

Page 3

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro, and K₁ is Asn; then one or more A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

26. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-J₁-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu;

J₁ is Ser, Pro, Leu, Ile or Thr;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy, and provided that

Appl. No. 10/643,681

Docket No.: 18528.642/ 254/057CON

Page 4

when

(a) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Pro and K₁ is Asn; or

(b) A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Val, J₁ is Ser and K₁ is Asn;

then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

27. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Leu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-G₁-Asn-H₁-Gly-²⁵I₁-J₁-Leu-Pro-Pro-³⁰Thr-K₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu;

K₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that

Appl. No. 10/643,681

Docket No.: 18528.642/ 254/057CON

Page 5

when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val and K₁ is Asn; then one or more of A₁ to K₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

28. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:

¹A₁-X-Asn-Thr-⁵Ala-Thr-Y-Ala-Thr-¹⁰Gln-Arg-Ileu-B₁-Asn-¹⁵Phe-Leu-C₁-D₁-E₁-²⁰F₁-
G₁-Asn-H₁-Gly-²⁵Pro-I₁-Leu-Pro-Pro-³⁰Thr-J₁-Val-Gly-Ser-³⁵Asn-Thr-Tyr-Z

wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala, Ser or Thr;

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr;

I₁ is Ile, Val, Ala or Leu

J₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and

provided that when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Val and J₁ is Asn; then one or more of A₁ to J₁ is a D-amino acid and Z is selected from the group consisting of alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy.

Appl. No. 10/643,681

Docket No.: 18528.642/ 254/057CON

Page 6

29. (Previously presented) The method of claim 24 wherein said amylin agonist is any one of $^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$, $\text{des-}^1\text{Lys}^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$, $^{25,28,29}\text{Pro-h-amylin}$, $\text{des-}^1\text{Lys}^{25,28,29}\text{Pro-h-amylin}$, $^{18}\text{Arg}^{25,28,29}\text{Pro-h-amylin}$, $\text{des-}^1\text{Lys}^{18}\text{Arg}^{25,28,29}\text{Pro-h-amylin}$, $^{25}\text{Pro}^{26}\text{Val}^{28,29}\text{Pro-h-amylin}$, or $\text{des-}^1\text{Lys}^{25}\text{Pro}^{26}\text{Val}^{28,29}\text{Pro-h-amylin}$.

30. (Previously presented) The method of claim 24 wherein the amylin agonist is $^{25,28,29}\text{Pro-h-amylin}$.

31-37. (Canceled)

38. (Previously presented) The method of claim 24 wherein the mammal has diabetes.

39. (Previously presented) The method of claim 38 wherein the diabetes is type 1.

40. (Previously presented) The method of claim 38 wherein the diabetes is type 2.

41. (Previously presented) The method of claim 25 wherein the mammal has diabetes.

42. (Previously presented) The method of claim 41 wherein the diabetes is type 1.

43. (Previously presented) The method of claim 41 wherein the diabetes is type 2.

44. (Previously presented) The method of claim 26 wherein the mammal has diabetes.

45. (Previously presented) The method of claim 44 wherein the diabetes is type 1.

46. (Previously presented) The method of claim 44 wherein the diabetes is type 2.

Appl. No. 10/643,681

Docket No.: 18528.642/ 254/057CON

Page 7

47. (Previously presented) The method of claim 27 wherein the mammal has diabetes.
48. (Previously presented) The method of claim 47 wherein the diabetes is type 1.
49. (Previously presented) The method of claim 47 wherein the diabetes is type 2.
50. (Previously presented) The method of claim 28 wherein the mammal has diabetes.
51. (Previously presented) The method of claim 50 wherein the diabetes is type 1.
52. (Previously presented) The method of claim 50 wherein the diabetes is type 2.
53. (Previously presented) The method of claim 30 wherein the mammal has diabetes.
54. (Previously presented) The method of claim 53 wherein the diabetes is type 1.
55. (Previously presented) The method of claim 53 wherein the diabetes is type 2.
56. (Previously presented) The method of claim 24 wherein the amylin agonist is an amylin agonist analogue having the following amino acid sequence:
 $^1\text{A}_1\text{-X-Asn-Thr-}^5\text{Ala-Thr-X-Ala-Thr-}^{10}\text{Gln-Arg-Leu-B}_1\text{-Asn-}^{15}\text{Phe-Leu-C}_1\text{-D}_1\text{-E}_1\text{-}^{20}\text{F}_1\text{-G}_1\text{-Asn-H}_1\text{-Gly-}^{25}\text{I}_1\text{-J}_1\text{-Leu-K}_1\text{-L}_1\text{-}^{30}\text{Thr-M}_1\text{-Val-Gly-Ser-}^{35}\text{Asn-Thr-Tyr-Z}$
wherein
A₁ is Lys, Ala, Ser, Hydrogen or acetylated Lys;
B₁ is Ala, Ser or Thr;

Appl. No. 10/643,681

Docket No.: 18528.642/ 254/057CON

Page 8

C₁ is Val, Leu or Ile;

D₁ is His or Arg;

E₁ is Ser or Thr;

F₁ is Ser, Thr, Gln or Asn;

G₁ is Asn, Gln or His;

H₁ is Phe, Leu or Tyr,

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu;

K₁ is Ser, Pro, Leu, Ile or Thr;

L₁ is Ser, Pro or Thr;

M₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage, wherein said intramolecular linkage comprises a disulfide bond, a lactam or a thioether linkage; and Z is an amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; and provided that

(a) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Phe, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Ser, and M₁ is Asn;

(b) when A₁ is Lys, B₁ is Ala, C₁ is Ile, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn;

(c) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Thr, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn;

(d) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Pro, L₁ is Pro, and M₁ is Asn;

(e) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Ser, L₁ is Pro and M₁ is Asn; or

(f) when A₁ is Lys, B₁ is Thr, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is His, H₁ is Leu, I₁ is Ala, J₁ is Ala, K₁ is Leu, L₁ is Pro and M₁ is Asp;

then one or more of any of A₁ to M₁ is not an L-amino acid and Z is not amino.

Appl. No. 10/643,681

Docket No.: 18528.642/ 254/057CON

Page 9

57. (Previously presented) The method of claim 56 wherein the mammal has diabetes.
58. (Previously presented) The method of claim 57 wherein the diabetes is type 1.
59. (Previously presented) The method of claim 57 wherein the diabetes is type 2.
- 60-69. (Canceled)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.